


**Amendments to the Claims:**


This listing of claims will replace all prior versions, and listings, of claims in the application:

**Listing of Claims:**

1. (currently amended) A process for preparing a compound of formula (I)



where  $R^4$  and  $R^5$  are independently selected from hydrogen, halo, nitro, cyano, hydroxy, fluoromethyl, difluoromethyl, trifluoromethyl, trifluoromethoxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, ureido,  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{1-6}$ alkoxy,  $C_{1-6}$ alkanoyl,  $C_{1-6}$ alkanoyloxy,  $N$ -( $C_{1-6}$ alkyl)amino,  $N,N$ -( $C_{1-6}$ alkyl)<sub>2</sub>amino,  $C_{1-6}$ alkaneylamino,  $N$ -( $C_{1-6}$ alkyl)carbamoyl,  $N,N$ -( $C_{1-6}$ alkyl)<sub>2</sub>carbamoyl,  $C_{1-6}$ alkylS(O)<sub>a</sub>, wherein  $a$  is 0 to 2,  $C_{1-6}$ alkoxy-carbonyl,  $C_{1-6}$ alkoxy-carbonylamino,  $N$ -( $C_{1-6}$ alkyl)sulphamoyl,  $N,N$ -( $C_{1-6}$ alkyl)<sub>2</sub>sulphamoyl,  $C_{1-6}$ alkylsulphenylamino and  $C_{1-6}$ alkylsulphenyl- $N$ -( $C_{1-6}$ alkyl)amino; and  $R^6$  is hydrogen or a protecting group, which process comprises cyclisation of a compound of formula (II)



where  $R^4$ ,  $R^5$  and  $R^6$  are as defined in relation to formula (I), and  $R^7$  is a nitrogen protecting group; and  
removing the group  $R^7$ ;  
and thereafter optionally removing any protecting group  $R^6$ .


2. (previously presented) A process according to claim 1 wherein  $R^7$  is a group of sub-formula (i)



where R<sup>8</sup> is a straight chain alkyl group of from 1 to 6 carbon atoms.

3-11 (cancelled)

12. (previously presented) A process according to claim 1, for preparing a compound of formula (I) where R<sup>6</sup> is hydrogen, wherein the process further comprises the step of reacting the compound of formula (I) obtained with an amine of formula (XIII)



where R<sup>14</sup> is selected from hydrogen or C<sub>1-6</sub>alkyl,

m is an integer of from 0 to 4,

each R<sup>15</sup> is the same or different and is selected from hydrogen, halo, nitro, cyano, hydroxy,

amino, carboxy, carbamoyl, mercapto, sulphamoyl, ureido, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl,

C<sub>2-6</sub>alkynyl, C<sub>1-6</sub>alkoxy, C<sub>1-6</sub>alkanoyl, C<sub>1-6</sub>alkanoyloxy, *N*-(C<sub>1-6</sub>alkyl)amino,

*N,N*-(C<sub>1-6</sub>alkyl)<sub>2</sub>amino, C<sub>1-6</sub>alkanoylamino, *N*-(C<sub>1-6</sub>alkyl)carbamoyl,

*N,N*-(C<sub>1-6</sub>alkyl)<sub>2</sub>carbamoyl, C<sub>1-6</sub>alkylS(O)<sub>a</sub> wherein a is 0 to 2, C<sub>1-6</sub>alkoxycarbonyl,

C<sub>1-6</sub>alkoxycarbonylamino, *N*-(C<sub>1-6</sub>alkyl)sulphamoyl, *N,N*-(C<sub>1-6</sub>alkyl)<sub>2</sub>sulphamoyl,

C<sub>1-6</sub>alkylsulphonylamino, C<sub>1-6</sub>alkylsulphonyl-*N*-(C<sub>1-6</sub>alkyl)amino, C<sub>3-6</sub>cycloalkyl,

C<sub>3-6</sub>cycloalkylC<sub>1-6</sub>alkyl, aryl, arylC<sub>1-6</sub>alkyl, heterocyclic group and (heterocyclic

group)C<sub>1-6</sub>alkyl; wherein R<sup>15</sup> may be optionally substituted on carbon by one or more

groups selected from P and wherein if said heterocyclic group contains an -NH- moiety

that nitrogen may be optionally substituted by a group selected from R;

each R<sup>16</sup> is the same or different and is selected from hydrogen or C<sub>1-6</sub>alkyl;

R<sup>17</sup> is selected from hydrogen, halo, nitro, cyano, hydroxy, fluoromethyl, difluoromethyl,

trifluoromethyl, trifluoromethoxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl,

ureido, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>1-6</sub>alkoxy, C<sub>1-6</sub>alkanoyl, C<sub>1-6</sub>alkanoyloxy,

*N*-(C<sub>1-6</sub>alkyl)amino, *N,N*-(C<sub>1-6</sub>alkyl)<sub>2</sub>amino, C<sub>1-6</sub>alkanoylamino, *N*-(C<sub>1-6</sub>alkyl)carbamoyl,

*N,N*-(C<sub>1-6</sub>alkyl)<sub>2</sub>carbamoyl, *N*-(C<sub>1-6</sub>alkyl)-*N*-(C<sub>1-6</sub>alkoxy)carbamoyl, C<sub>1-6</sub>alkylS(O)<sub>a</sub> wherein

a is 0 to 2, C<sub>1-6</sub>alkoxycarbonyl, C<sub>1-6</sub>alkoxycarbonylamino, *N*-(C<sub>1-6</sub>alkyl)sulphamoyl,

*N,N*-(C<sub>1-6</sub>alkyl)<sub>2</sub>sulphamoyl, sulphamoylamino, *N*-(C<sub>1-6</sub>alkyl)sulphamoylamino,

*N,N*-(C<sub>1-6</sub>alkyl)<sub>2</sub>sulphamoylamino, C<sub>1-6</sub>alkylsulphonylamino,

C<sub>1-6</sub>alkylsulphonylamino, C<sub>1-6</sub>alkylsulphonyl-*N*-(C<sub>1-6</sub>alkyl)amino and a group -E-F-G-H;

wherein E and G are independently selected from a direct bond, -O-, -S-, -SO-, -SO<sub>2</sub>-, -OC(O)-, -C(O)O-, -C(O)-, -NR<sup>a</sup>-, -NR<sup>a</sup>C(O)-, -C(O)NR<sup>a</sup>-, -SO<sub>2</sub>NR<sup>a</sup>-, -NR<sup>a</sup>SO<sub>2</sub>-, -NR<sup>a</sup>C(O)NR<sup>b</sup>-, -OC(O)NR<sup>a</sup>-, -NR<sup>a</sup>C(O)O-, -NR<sup>a</sup>SO<sub>2</sub>NR<sup>b</sup>-, -SO<sub>2</sub>NR<sup>a</sup>C(O)- and -C(O)NR<sup>a</sup>SO<sub>2</sub>-; wherein R<sup>a</sup> and R<sup>b</sup> are independently selected from hydrogen or C<sub>1-6</sub>alkyl which is optionally substituted by a group V;

F is C<sub>1-6</sub>alkylene optionally substituted by one or more Q or a direct bond;

H is selected from aryl, C<sub>3-6</sub>cycloalkyl and heterocyclic groups; wherein H may be optionally substituted on carbon by one or more groups selected from S and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from T;


P, S and Q are independently selected from halo, nitro, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, ureido, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>1-6</sub>alkoxy, C<sub>1-6</sub>alkanoyl, C<sub>1-6</sub>alkanoyloxy, *N*-(C<sub>1-6</sub>alkyl)amino, *N,N*-(C<sub>1-6</sub>alkyl)<sub>2</sub>amino, C<sub>1-6</sub>alkanoylamino, *N*-(C<sub>1-6</sub>alkyl)carbamoyl, *N,N*-(C<sub>1-6</sub>alkyl)<sub>2</sub>carbamoyl, *N*-(C<sub>1-6</sub>alkyl)-*N*-(C<sub>1-6</sub>alkoxy)carbamoyl, C<sub>1-6</sub>alkylS(O)<sub>a</sub> wherein a is 0 to 2, C<sub>1-6</sub>alkoxycarbonyl, C<sub>1-6</sub>alkoxycarbonylamino, *N*-(C<sub>1-6</sub>alkyl)sulphamoyl, *N,N*-(C<sub>1-6</sub>alkyl)<sub>2</sub>sulphamoyl, C<sub>1-6</sub>alkylsulphonylamino, C<sub>1-6</sub>alkylsulphonyl-*N*-(C<sub>1-6</sub>alkyl)amino, C<sub>3-6</sub>cycloalkyl, aryl and heterocyclic group; wherein P, S and Q may be optionally and independently substituted on carbon by one or more groups selected from V and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from U;

V is selected from halo, nitro, cyano, hydroxy, trifluoromethoxy, trifluoromethyl, amino, carboxy, carbamoyl, mercapto, sulphamoyl, methyl, ethyl, methoxy, ethoxy, acetyl, acetoxo, methylamino, ethylamino, dimethylamino, diethylamino, *N*-methyl-*N*-ethylamino, acetylamino, *N*-methylcarbamoyl, *N*-ethylcarbamoyl, *N,N*-dimethylcarbamoyl, *N,N*-diethylcarbamoyl, *N*-methyl-*N*-ethylcarbamoyl, methylthio, ethylthio, methylsulphinyl, ethylsulphinyl, mesyl, ethylsulphonyl, methoxycarbonyl, ethoxycarbonyl, *N*-methylsulphamoyl, *N*-ethylsulphamoyl, *N,N*-dimethylsulphamoyl, *N,N*-diethylsulphamoyl, *N*-methyl-*N*-ethylsulphamoyl, morpholino, morpholinocarbonyl, *N*-benzylcarbamoyl, and 4-hydroxypiperidinocarbonyl;

R, T and U are independently selected from C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkanoyl, C<sub>1-4</sub>alkylsulphonyl, C<sub>1-4</sub>alkoxycarbonyl, carbamoyl, *N*-(C<sub>1-4</sub>alkyl)carbamoyl, *N,N*-(C<sub>1-4</sub>alkyl)carbamoyl, phenyl,

benzyl, benzyloxycarbonyl, benzoyl and phenylsulphonyl wherein R, T and U may be optionally and independently substituted on carbon by one or more groups selected from V;

producing a compound of formula (XIV)



where R<sup>4</sup>, R<sup>5</sup>, R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup> and m are as defined above,

or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof.

13-15. (cancelled)